

Binary Response Models

1 Introduction

Many dependent variables of interest in economics and other social sciences can only take two values. The two possible outcomes are usually denoted by 0 and 1. Such variables are called *dummy* variables or *dichotomous* variables. Some examples:

- The labor market status of a person. The variable takes the value 1 if a person is employed and 0 if he is unemployed. The values 1 and 0 can be assigned arbitrarily.
- Voting behavior of a person. The variable takes 1 if the person votes in favor of a new policy and 0 otherwise. Again the values 1 and 0 are arbitrary.

The expected value of a dichotomous variable $y_i \in \{0, 1\}$ is the probability that it takes the value 1:

$$E(y_i) = 0 \cdot P(y_i = 0) + 1 \cdot P(y_i = 1) = P(y_i = 1).$$

The linear regression model,

$$y_i = x_i' \beta + u_i, \quad E(u_i | x_i) = 0$$

is called *linear probability model* in this context. This linear model is not an adequate statistical model as the expected value $E(y_i | x_i) = x_i' \beta$ can lie outside $[0, 1]$ and does not represent a probability. In addition, the error term is heteroscedastic as $V(u_i | x_i) = x_i' \beta (1 - x_i' \beta)$ depends on x_i .

2 The Econometric Model: Probit and Logit

Binary response models directly describe the response probabilities $P(y_i = 1)$ of the dependent variable y_i .

Consider a sample of N independently and identically distributed (i.i.d.) observations $i = 1, \dots, N$ of the dependent dummy variable y_i and a $(K+1)$ -dimensional vector x'_i of explanatory variables including a constant. The probability that the dependent variable takes value 1 is modeled as

$$P(y_i = 1|x_i) = F(z_i) = F(x'_i\beta)$$

where β is a $(K + 1)$ -dimensional column vector of parameters and

$$z_i = x'_i\beta$$

is a *single linear index*. The *transformation function* F maps the single index into $[0,1]$ and satisfies in general

$$F(-\infty) = 0, \quad F(\infty) = 1, \quad \partial F(z)/\partial z > 0.$$

The *probit* model assumes that the transformation function F is the cumulative density function (cdf) of the standard normal distribution. The response probabilities are then

$$P(y_i = 1|x_i) = \Phi(x'_i\beta) = \int_{-\infty}^{x'_i\beta} \phi(t) dt = \int_{-\infty}^{x'_i\beta} \frac{1}{\sqrt{2\pi}} e^{-\frac{1}{2}t^2} dt$$

where $\phi(\cdot)$ is the pdf and $\Phi(\cdot)$ the cdf of the standard normal distribution.

In the *logit* model, the transformation function F is the logistic function. The response probabilities are then

$$P(y_i = 1|x_i) = \frac{e^{x'_i\beta}}{1 + e^{x'_i\beta}} = \frac{1}{1 + e^{-x'_i\beta}}$$

Figure 1 shows the transformation function F for the two models.

Note: The Logit and Probit model are almost identical and the choice of the model is usually arbitrary. However, the parameters β of the two




Figure 1: The transformation function in the probit and logit model.

models are scaled differently. Multiplying the parameters in the probit model by 1.6 are approximately the same as the logit estimates.¹

3 Latent Variable Model

There is an alternative interpretation that gives rise to the probit (and analogously the logit) model. Consider a *latent* variable which is not observed by the researcher and linearly depends on x_i

$$y_i^* = x_i' \beta + u_i, \quad E(u_i) = 0$$

The latent variable y_i^* can be interpreted as the utility difference between choosing $y_i = 1$ and 0. In is then called a *random utility model*.

¹The factor 1.6 equals the first derivative of F at $x_i' \beta = 0$ and is in most applications the appropriate rescaling. A different approach is to equal the standard deviation of the distribution for which F is the cdf. For the probit model the standard deviation is 1 and for the logit model $\pi/\sqrt{3} \cong 1.81$.




Figure 2: The probit model with a latent variable. $N = 30$, $K = 2$ (a constant and one explanatory variable) and $\beta = (-2, 0.5)$.

Only the choice y_i is observed by the researcher. An individual chooses $y_i = 1$ if the latent variable is positive and 0 otherwise, hence the observed variable is

$$y_i = \begin{cases} 1 & \text{if } y_i^* > 0 \\ 0 & \text{if } y_i^* \leq 0 \end{cases}$$

Furthermore, assume that the individual observations (x_i, y_i) are i.i.d., that the explanatory variables are *exogenous* and that the error term is

normally distributed and homoskedastic

$$u_i|x_i \sim N(0, \sigma^2)$$

The probability that individual i chooses $y_i = 1$ can now be derived from the latent variable and the decision rule, i.e.

$$\begin{aligned} P(y_i = 1|x_i) &= P(y_i^* > 0|x_i) = P(x_i'\beta + u_i > 0|x_i) = P(u_i > -x_i'\beta|x_i) \\ &= 1 - \Phi(-x_i'\beta/\sigma) = \Phi(x_i'\beta/\sigma). \end{aligned}$$

The probit model arises when σ^2 is set to unity.

Note: β_k and σ are *not separately identified* as only the ratio β_k/σ can be estimated. Figure 2 visualizes the latent variable model.

4 Interpretation of the Parameters

Different from the linear regression model, the parameters β cannot directly be interpreted as marginal effects on the dependent variable y_i . In some situations, the index function $z_i = x_i'\beta$ has a clear interpretation in a theoretical model and the marginal effect β_k of a change in the independent variable x_{ik} on y_i^* is meaningful. Even then, the marginal effect is only identified if there is reason to set σ^2 to unity.

In general, we are interested in the marginal effect of a change in x_{ik} on the expected value of the observed variable y_i , i.e.

$$\begin{aligned} \text{Probit:} \quad & \frac{\partial E(y_i|x_i)}{\partial x_{ik}} = \frac{\partial P(y_i = 1|x_i)}{\partial x_{ik}} = \phi(x_i'\beta) \beta_k \\ \text{Logit:} \quad & \frac{\partial E(y_i|x_i)}{\partial x_{ik}} = \frac{\partial P(y_i = 1|x_i)}{\partial x_{ik}} = \frac{e^{x_i'\beta}}{(1 + e^{x_i'\beta})^2} \beta_k \end{aligned}$$

This marginal effect depends on the characteristics of all x_{ik} for observation i . Therefore, any individual has a different marginal effect. There are several ways to summarize and report the information in the model. A first possibility is to present the marginal effects for the “mean type”,

i.e. $x_i = \bar{x}_i$, the “median type”, or some interesting extreme types. A second approach is to calculate the marginal effects for all observations in the sample and report the mean of the effects.

The estimated model can also be used for predictions

$$\begin{aligned} \text{Probit: } \hat{P}(y_i = 1|x_i) &= \Phi(x_i' \hat{\beta}) \\ \text{Logit: } \hat{P}(y_i = 1|x_i) &= \frac{e^{x_i' \hat{\beta}}}{1 + e^{x_i' \hat{\beta}}} \end{aligned}$$

This information can be aggregated to, for example, the predicted number of observations with $y_i = 1$. There are two prediction methods for this aggregate: (1) assume $\hat{y}_i = 1$ if $\hat{P}_i > 0.5$ and calculate $\sum_i \hat{y}_i$ or (2) sum the predicted choice probabilities $\sum_i \hat{P}(y_i = 1|x_i)$. The two measures can be contrasted to the actual numbers. Method 1 also allows to compare actual and predicted outcomes for any observation. It is also often interesting to report and contrast predicted numbers for certain types of individuals.

5 Estimation with Maximum Likelihood

The probit and logit models are estimated by maximum likelihood (ML). Assuming independence across observations, the likelihood function is

$$\begin{aligned} \mathcal{L} &= \prod_{\{i|y_i=0\}} P(y_i = 0|x_i) \prod_{\{i|y_i=1\}} P(y_i = 1|x_i) \\ &= \prod_{i=1}^N [1 - F(z_i)]^{1-y_i} F(z_i)^{y_i} \end{aligned}$$

where $P(y_i = 1|x_i) = F(z_i) = \Phi(z_i)$ in the probit model and $P(y_i = 1|x_i) = F(z_i) = e^{z_i}/(1 + e^{z_i})$ in the logit model. The corresponding log likelihood function is

$$\log \mathcal{L} = \sum_{i=1}^N [y_i \log F(z_i) + (1 - y_i) \log (1 - F(z_i))]$$

The first order conditions for an optimum are in general, for all k including a constant $x_{i0} = 1$

$$\frac{\partial \log \mathcal{L}}{\partial \beta_k} = \sum_{i=1}^N \left[y_i \frac{f(z_i)}{F(z_i)} + (1 - y_i) \frac{-f(z_i)}{1 - F(z_i)} \right] x_{ik} = 0$$

where $f(z) \equiv \partial F(z)/\partial z$. This simplifies in the probit model to

$$\frac{\partial \log \mathcal{L}}{\partial \beta_k} = \sum_{\{i|y_i=0\}} \frac{-\phi(z_i)}{1 - \Phi(z_i)} x_{ik} + \sum_{\{i|y_i=1\}} \frac{\phi(z_i)}{\Phi(z_i)} x_{ik} = 0$$

and in the logit model to

$$\frac{\partial \log \mathcal{L}}{\partial \beta_k} = \sum_{i=1}^N \left(y_i - \frac{e^{z_i}}{1 + e^{z_i}} \right) x_{ik} = 0.$$

There is no analytical solution to these FOCs and numerical optimization routines are used. The log likelihood function can be shown to be globally concave for both models and numerical routines converge well to the unique global maximum.

The ML estimator of β is consistent and asymptotically normally distributed. The approximate distribution in large samples is

$$\hat{\beta} \stackrel{A}{\sim} N(\beta, I(\beta)^{-1})$$

where the inverse information matrix I^{-1} is estimated by one of the standard ML procedures (inverse expected H, inverse Hessian, BHHH, or Eicker-Huber-White-Sandwich). Asymptotic hypothesis tests are performed as Wald, likelihood ratio or lagrange multiplier tests.

The ML estimation of the probit model (and analogously the logit model) rests on the strong assumption that the latent error term is normally distributed and homoscedastic. The ML estimator is inconsistent in the presence of heteroscedasticity and robust covariance estimators cannot solve this. Several *semi-parametric* estimation strategies have been proposed that relax the distributional assumption about the error term. See Horowitz and Savin (2001) for an introduction and Gerfin (1996) for a nice comparison of different estimators.

6 Estimation with OLS

Despite the logical inconsistency of the linear probability model, OLS can be used to estimate binary choice models. The estimated OLS slope coefficients are estimates for the average marginal effects of the true non-linear model. In practice, the OLS slope coefficients will be very similar to the average marginal effects calculated after probit or logit estimation. However, it is very important to report robust (Eicker-White) standard errors because of the intrinsic heteroscedasticity of the linear probability model.

The linear probability model has in practice several advantages over probit or logit estimation: it is easier to calculate, the parameters are directly interpretable, fixed effects and instrumental variables estimators can easily be implemented. Note that adding fixed effects as dummy variables in the probit or logit model will yield biased estimates.

7 Implementation in Stata 11.0

Stata estimates the probit model by the command

```
probit devar [indepvars]
```

and the logit model by

```
logit devar [indepvars]
```

where *devar* is the dependent dummy variable and *indepvars* is a list of explanatory variables. For example,

```
webuse auto.dta
probit foreign weight mpg
```

Stata reports the inverse hessian matrix as default covariance estimator. The option `vce(opg)` reports BHHH and option `vce(robust)` the sandwich covariance estimator.

Response probabilities are estimated for each observation with the post-estimation command `predict`. For example,

```
predict p_foreign, pr
```

predicts the probability of being foreign for each car model. Marginal effects for specific types are calculated with the post-estimation command `mfxf`. For example,

```
mfxf
```

calculates the marginal effects for the mean type, e.g. a car with average weight and mpg. The marginal effects for a specific type, e.g. a car with weight of 2000 lbs. and 40 mpg is reported by

```
mfxf, at(weight = 2000, mpg = 40)
```

In the case of explanatory dummy variables, Stata reports the exact discrete effect. Marginal effects for each individual in the sample can be calculated manually. For example in the probit model,

```
probit foreign weight mpg
predict xb, xb
predictnl mfx_weight = normalden(xb)*_b[weight]
```

or in the logit model

```
logit foreign weight mpg
predict p, pr
predictnl mfx_weight = p*(1-p)*_b[weight]
```

The post-estimation command `predictnl` cannot be used to calculate confidence intervals for individual marginal effects as the error as prediction error in p and xb , respectively, is ignored.

References

Introductory textbooks

- Stock, James H. and Mark W. Watson (2007), *Introduction to Econometrics*, 2nd ed., Pearson Addison-Wesley. Chapter 11.
- Wooldridge, Jeffrey M. (2009), *Introductory Econometrics: A Modern Approach*, 4th ed., South-Western Cengage Learning. Chapters 17.1.
- Aldrich, John and Forrest D. Nelson (1984), *Linear Probability, Logit and Probit Models*, Sage University Press.

Advanced textbooks

- Cameron, A. Colin and Pravin K. Trivedi (2005), *Microeconometrics: Methods and Applications*, Cambridge University Press. Chapter 14.
- Wooldridge, Jeffrey M. (2002), *Econometric Analysis of Cross Section and Panel Data*, MIT Press. Chapter 15.
- Maddala, G.S. (1983), *Limited-Dependent and Qualitative Variables in Econometrics*, Cambridge: Cambridge University Press. Chapter 2.

Articles

- Gerfin Michael (1996), Parametric and Semi-Parametric Estimation of the Binary Response Model of Labour Market Participation, *Journal of Applied Econometrics*, 11, 321-339.
- Horowitz, Joel and N. Savin (2001), Binary Response Models: Logits, Probits and Semiparametrics, *Journal of Economic Perspectives*, 15(4), 43-56.